

## Ask your doctor today about Latisse™

### Can anyone use Latisse™?

You should not use Latisse™ solution if you are allergic to one of its ingredients. Also, tell your doctor or eye care specialist if any of the following apply to you:

- **Previously diagnosed with eye pressure problems:** People with a history of eye pressure problems (specifically *elevated intraocular pressure*) should speak to an eye care specialist before starting Latisse™
- **Currently taking medication for eye pressure problems:** People taking medication to lower their IOP should be closely monitored while using Latisse™ solution
- **Have risk factors for glaucoma:** People with risk factors for an eye condition called *glaucoma*, including a family history of *glaucoma*, should speak to an eye care specialist before starting Latisse™



### Is Latisse™ safe?

The safety and effectiveness of Latisse™ was shown in a clinical trial. Ask your doctor for more information about the safety of Latisse™ solution, and see the Important Safety Information on the back cover of this brochure.



### What are the possible side effects with Latisse™?

The most common side effects after using Latisse™ are eye itching or eye redness. These were reported by about 4% of users in clinical trial.<sup>1</sup> Latisse™ solution may cause other less common side effects, which typically occur on the skin close to where Latisse™ is applied, or in the eyes. These include skin darkening, eye irritation, dryness, or redness of the eyelids.<sup>1</sup>

If you develop a new eye condition (like an eye trauma or infection), experience a sudden decrease in vision, have eye surgery, or develop any eye reactions, particularly an infection and eyelid reactions, you should immediately talk to your doctor about whether or not to continue using Latisse™ solution.

Please see Important Safety Information on the back cover.

**Latisse™**  
(bimatoprost ophthalmic solution) 0.03%